

MİKRODALGA 2

GEREKLİ OLABİLECEK FORMÜLLER:

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} \cosh \gamma l & Z_c \sinh \gamma l \\ \frac{1}{Z_c} \sinh \gamma l & \cosh \gamma l \end{bmatrix}$$

$$\Gamma_{in} = S_{11} + \frac{S_{12}S_{21}\Gamma_L}{1 - S_{22}\Gamma_L}$$

$$\Gamma_{out} = S_{22} + \frac{S_{12}S_{21}\Gamma_S}{1 - \Gamma_S S_{11}}$$

$$\Gamma_{in} = \frac{A + \frac{B}{Z_{02}} - Z_{01}C - D \frac{Z_{01}}{Z_{02}}}{A + \frac{B}{Z_{02}} + Z_{01}C + D \frac{Z_{01}}{Z_{02}}}$$

$$T_{21} = \frac{2}{A + \frac{B}{Z_{02}} + Z_{01}C + D \frac{Z_{01}}{Z_{02}}}$$

$$k = \frac{\sqrt{\frac{Z_{oe}}{Z_{oo}}} - \sqrt{\frac{Z_{oo}}{Z_{oe}}}}{\sqrt{\frac{Z_{oe}}{Z_{oo}}} + \sqrt{\frac{Z_{oo}}{Z_{oe}}}}$$

$$k = \frac{\rho - 1}{\rho + 1}$$

$$\rho = \frac{1+k}{1-k}$$

$$K = \frac{j k \sin \theta}{\sqrt{1 - k^2} \cos \theta + j \sin \theta} = \frac{V_3^-}{V_1^+}$$

$$G = \frac{\sqrt{1 - k^2}}{\sqrt{1 - k^2} \cos \theta + j \sin \theta} = \frac{V_4^-}{V_1^+}$$

$$\frac{K^2}{K^2 + G^2}$$

$$S_{11} = \frac{A + \frac{B}{Z_0} - Z_0 C - D}{A + \frac{B}{Z_0} + Z_0 C + D}$$

$$S_{21} = \frac{2}{A + \frac{B}{Z_0} + Z_0 C + D}$$

$$Z = R + jX$$

$$|\Gamma_{out}| = \left| \frac{S_{22} + S_{12}S_{21}\Gamma_S}{1 - S_{11}\Gamma_S} \right|$$

$$|\Gamma_{in}| = \left| \frac{S_{11} + S_{12}S_{21}\Gamma_L}{1 - \Gamma_L S_{22}} \right|$$

$$K = \frac{1 + |\Delta|^2 - |S_{11}|^2 - |S_{22}|^2}{2 - |S_{12}||S_{21}|}$$

$$\Delta = S_{11}S_{22} - S_{12}S_{21}$$

$$G_P = \frac{P_L}{P_{IN}}$$

$$G_T = \frac{P_L}{P_{AVS}}$$

$$G_A = \frac{P_{AVN}}{P_{AVS}}$$

$$G_T = \frac{|S_{21}|^2(1 - |\Gamma_S|^2)(1 - |\Gamma_L|^2)}{|(1 - \Gamma_S S_{11})(1 - \Gamma_L S_{22}) - \Gamma_S \Gamma_L S_{12} S_{21}|^2}$$

$$G_T = \frac{|S_{21}|^2(1 - |\Gamma_S|^2)(1 - |\Gamma_L|^2)}{|1 - S_{11}\Gamma_S|^2|1 - \Gamma_{out}\Gamma_L|^2}$$

$$G_T = \frac{|S_{21}|^2(1 - |\Gamma_S|^2)(1 - |\Gamma_L|^2)}{|1 - \Gamma_{in}\Gamma_S|^2|1 - S_{22}\Gamma_L|^2}$$

$$S'_{11} = S'_{22} = 0 \quad , \quad G_{Amaks} = \frac{|S_{21}|}{|S_{12}|} (K - \sqrt{K^2 - 1})$$

$$G_A = \frac{|S_{21}|^2(1 - |\Gamma_S|^2)}{|1 - S_{11}\Gamma_S|^2|1 - |\Gamma_{out}|^2|}$$

$$G_p = \frac{|S_{21}|^2(1 - |\Gamma_L|^2)}{|1 - |\Gamma_{in}|^2||1 - S_{22}\Gamma_L|^2}$$

$$G_{TU} = \frac{1 - |\Gamma_S|^2}{|1 - S_{11}\Gamma_S|^2} |S_{21}|^2 \frac{1 - |\Gamma_L|^2}{|1 - S_{22}\Gamma_L|^2}$$

$$G_S = \frac{1 - |\Gamma_S|^2}{|1 - S_{11}\Gamma_S|^2}$$

$$G_L = \frac{1 - |\Gamma_L|^2}{|1 - S_{22}\Gamma_L|^2}$$

$$G_O = |S_{21}|^2$$

$$G_{Smax} = \frac{1}{1 - |S_{11}|^2}$$

$$G_{Lmax} = \frac{1}{1 - |S_{22}|^2}$$

$$G_{TUm\max} = \frac{1}{1 - |S_{11}|^2} |S_{21}|^2 \frac{1}{1 - |S_{22}|^2}$$